


VIOLENT DEATH IN NORTH CAROLINA


The North Carolina Violent Death Reporting System (NC-VDRS) is a CDC-funded state-wide surveillance system that collects detailed information on deaths that occur in N.C. resulting from violence: homicide, suicide, unintentional firearm deaths, legal intervention, and deaths for which intent could not be determined. NC-VDRS is a multi-source system that gathers information from death certificates, medical examiner reports, and law enforcement reports. The goal of this system is to aid researchers, legislators, and community interest groups in the development of public health prevention strategies to reduce violent deaths. NC-VDRS began collecting data in January, 2004. This document summarizes all deaths from violence among N.C. residents for 2006.



- In 2006, 1,785 N.C. residents died by violence.
- There were 1,106 suicides (62%), 599 homicides (34%), 14 unintentional firearm deaths (1%), 24 deaths from legal interventions (1%), and 42 deaths of undetermined intent (2%) in 2006.


- The pattern by sex was the same for both suicides and homicides; three-fourths (77%) of victims were male and one-fourth of victims were female.
- Patterns of suicides and homicides differed by race. Most suicide victims were white (90%), while 9% were black.
- In contrast, approximately 55% of homicide victims were black, and 40% were white.

Violent Deaths by Gender and Race: N.C., 2006




*Other includes American Indians, Asians, and other races.

Violent Death Rates by Age Group: N.C., 2006



- Suicides and homicides displayed different age patterns.
- Homicide rates peaked in the 20-24 year age group, with a smaller peak in infancy.
- Suicides peaked in midlife (45-54 years).

Method of Death: N.C., 2006



- The majority of homicides (69%) and suicides (59%) were committed using firearms.
- The second most common method of homicide was sharp instruments (12%).
- The second most common method of suicide was poisoning (19%).
- The third most common method of death for homicides and suicides was unarmed assault (6%) and hanging/suffocation (16%), respectively.

*Other includes fire/burns, unarmed assault, blunt trauma, falls, and other methods of death.

The North Carolina Violent Death Reporting System is supported by Cooperative Agreement U17/CE423098-06 from the Centers for Disease Control and Prevention (CDC).

